

## **Product datasheet for TA371824S**

## **HDAC4 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human thyroid cancer

Predicted cell location: Cytoplasm and Nucleus

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

Immunogen:Synthetic peptide of human HDAC4Formulation:pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Gene Name: histone deacetylase 4

Database Link: Entrez Gene 9759 Human

P56524

**Background:** Histones play a critical role in transcriptional regulation, cell cycle progression, and

developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a

multiprotein complex with RbAp48 and HDAC3.

Synonyms: HA6116; HD4; HDAC-A; HDACA; KIAA0288




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA371824] (HDAC4 Antibody) at dilution 1/20 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using [TA371824] (HDAC4 Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)