

## **Product datasheet for TA371951**

# alpha 1 Spectrin (SPTA1) Rabbit Polyclonal Antibody

### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: Mouse heart tissue lysate

IHC: 60-450

Positive control: Human gastric cancer Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human SPTA1

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year
Predicted Protein Size: 280 kDa

**Gene Name:** spectrin alpha, erythrocytic 1

Database Link: Entrez Gene 6708 Human

P02549

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






#### Background:

Spectrin is an actin crosslinking and molecular scaffold protein that links the plasma membrane to the actin cytoskeleton, and functions in the determination of cell shape, arrangement of transmembrane proteins, and organization of organelles. It is a tetramer made up of alpha-beta dimers linked in a head-to-head arrangement. This gene is one member of a family of alpha-spectrin genes. The encoded protein is primarily composed of 22 spectrin repeats which are involved in dimer formation. It forms weaker tetramer interactions than non-erythrocytic alpha spectrin, which may increase the plasma membrane elasticity and deformability of red blood cells. Mutations in this gene result in a variety of hereditary red blood cell disorders, including elliptocytosis type 2, pyropoikilocytosis, and spherocytic hemolytic anemia.

Synonyms: EL2; HPP; HS3; SPH3; SPTA

# **Product images:**



Gel: 6%SDS-PAGE Lysate: 40 μg

Lane: Mouse heart tissue lysate


Primary antibody: TA371951 (SPTA1 Antibody) at

dilution 1/550

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 20 seconds



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using TA371951 (SPTA1 Antibody) at dilution 1/80 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using TA371951 (SPTA1 Antibody) at dilution 1/80, treated with synthetic peptide. (Original magnification: ×200)