

Product datasheet for TA372039S

WHSC1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human colorectal cancer

Predicted cell location: Nucleus

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human NSD2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Gene Name: Wolf-Hirschhorn syndrome candidate 1

Database Link: Entrez Gene 7468 Human

<u>096028</u>

Background: This gene encodes a protein that contains four domains present in other developmental

proteins: a PWWP domain, an HMG box, a SET domain, and a PHD-type zinc finger. It is expressed ubiquitously in early development. Wolf-Hirschhorn syndrome (WHS) is a malformation syndrome associated with a hemizygous deletion of the distal short arm of chromosome 4. This gene maps to the 165 kb WHS critical region and has also been involved in the chromosomal translocation t(4;14)(p16.3;q32.3) in multiple myelomas. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms. Some transcript variants are nonsense-mediated mRNA (NMD) decay candidates, hence not

represented as reference sequences. [provided by RefSeq, Jul 2008]



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

FLJ23286; KIAA1090; MGC176638; MMSET; NSD2; OTTHUMP00000196943; REIIBP; TRX5; WHS

Product images:



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA372039] (NSD2 Antibody) at dilution 1/40 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using [TA372039] (NSD2 Antibody) at dilution 1/40, treated with synthetic peptide. (Original magnification: ×200)