

## **Product datasheet for TA372136**

# **ATP5MJ Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: 293T and RAW264.7 cell Human fetal brain tissue Jurkat and 231

cell

Human heart tissue

Hela cell lysates

IHC: 40-200

Positive control: Human prostate cancer

Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Rabbit Host: Isotype: lgG

Clonality: Polyclonal

Synthetic peptide of human ATP5MPL Immunogen:

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year **Predicted Protein Size:** 7 kDa

Gene Name: chromosome 14 open reading frame 2

Database Link: Entrez Gene 9556 Human

P56378



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

Mitochondrial membrane ATP synthase (F1F0 ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F1 - containing the extramembraneous catalytic core and F0 - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F1 is coupled via a rotary mechanism of the central stalk subunits to proton translocation (Probable). Minor subunit required to maintain the ATP synthase population in the mitochondria (PubMed:24330338).

Synonyms: MP68; PLPM

### **Product images:**




Lysate: 40 µg Lane 1-7: 293T and RAW264.7 cell□Human fetal brain tissue□Jurkat and 231 cell□Human heart tissue□Hela cell lysates Primary antibody: TA372136 (ATP5MPL Antibody) at dilution 1/600 Secondary antibody: Goat anti rabbit IgG at

Gel: 12%SDS-PAGE


1/8000 dilution

Exposure time: 5 seconds




Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using TA372136 (ATP5MPL Antibody) at dilution 1/60 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using TA372136 (ATP5MPL Antibody) at dilution 1/60, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human brain tissue using TA372136 (ATP5MPL Antibody) at dilution 1/60 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human brain tissue using TA372136 (ATP5MPL Antibody) at dilution 1/60, treated with synthetic peptide. (Original magnification: ×200)