

Product datasheet for TA372469S

ELYS (AHCTF1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human ovarian cancer

Predicted cell location: Cytoplasm and Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human AHCTF1
Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

Gene Name: AT-hook containing transcription factor 1

Database Link: Entrez Gene 25909 Human

Q8WYP5

Background: Required for the assembly of a functional nuclear pore complex (NPC) on the surface of

chromosomes as nuclei form at the end of mitosis. May initiate NPC assembly by binding to chromatin and recruiting the Nup107-160 subcomplex of the NPC. Also required for the localization of the Nup107-160 subcomplex of the NPC to the kinetochore during mitosis and

for the completion of cytokinesis.

Synonyms: DKFZp434N093; ELYS; MST108; MSTP108; TMBS62




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:




Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA372469] (AHCTF1 Antibody) at dilution 1/20 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using [TA372469] (AHCTF1 Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA372469] (AHCTF1 Antibody) at dilution 1/20 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using [TA372469] (AHCTF1 Antibody) at dilution 1/20, treated with synthetic peptide. (Original magnification: ×200)