

Product datasheet for TA372696

GSTM2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 200-1000

WB positive control: Mouse skeletal muscle tissue lysate

IHC: 40-200

Positive control: Human ovarian cancer

Predicted cell location: Cytoplasm and Nucleus

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human GSTM2

Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 26 kDa

Gene Name: glutathione S-transferase mu 2 (muscle)

Database Link: Entrez Gene 2946 Human

P28161

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. This gene encodes a glutathione S-transferase that belongs to the mu class. The mu class of enzymes functions in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione. The genes encoding the mu class of enzymes are organized in a gene cluster on chromosome 1p13.3 and are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of certain drugs.

Synonyms:

GST4; GSTM; GSTM2-2; GTHMUS; MGC117303

Product images:



Gel: 12%SDS-PAGE Lysate: 40 µg


Lane: Mouse skeletal muscle tissue lysate Primary antibody: TA372696 (GSTM2 Antibody) at

dilution 1/300

Secondary antibody: Goat anti rabbit IgG at


1/8000 dilution

Exposure time: 2 minutes



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using TA372696 (GSTM2 Antibody) at dilution 1/30 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using TA372696 (GSTM2 Antibody) at dilution 1/30, treated with synthetic peptide. (Original magnification: ×200)