

# **Product datasheet for TA372821S**

### Junctional Adhesion Molecule C (JAM3) Rabbit Polyclonal Antibody

### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: K562 cell lysate

IHC: 40-200

Positive control: Human tonsil Predicted cell location: Cytoplasm

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human JAM3

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 35 kDa

**Gene Name:** junctional adhesion molecule 3

Database Link: Entrez Gene 83700 Human

Q9BX67



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

Tight junctions represent one mode of cell-to-cell adhesion in epithelial or endothelial cell sheets, forming continuous seals around cells and serving as a physical barrier to prevent solutes and water from passing freely through the paracellular space. The protein encoded by this immunoglobulin superfamily gene member is localized in the tight junctions between high endothelial cells. Unlike other proteins in this family, the this protein is unable to adhere to leukocyte cell lines and only forms weak homotypic interactions. The encoded protein is a member of the junctional adhesion molecule protein family and acts as a receptor for another member of this family. A mutation in an intron of this gene is associated with hemorrhagic destruction of the brain, subependymal calcification, and congenital cataracts. Alternative splicing results in multiple transcript variants.

Synonyms:

FLJ14529; JAM-2; JAM-3; JAM-C; JAMC

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: K562 cell lysate


Primary antibody: [TA372821] (JAM3 Antibody) at

dilution 1/1000

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 7 minutes



Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA372821] (JAM3 Antibody) at dilution 1/45 (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human tonsil tissue using [TA372821] (JAM3 Antibody) at dilution 1/45, treated with synthetic peptide. (Original magnification: ×200)