

## **Product datasheet for TA372837**

## **KCC4 (SLC12A7) Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC

Recommended Dilution: IHC: 25-100

Positive control: Human tonsil Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen:Synthetic peptide of human SLC12A7Formulation:pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year

**Gene Name:** solute carrier family 12 member 7

Database Link: Entrez Gene 10723 Human

Q9Y666

**Background:** Mediates electroneutral potassium-chloride cotransport when activated by cell swelling. May

mediate K+ uptake into Deiters' cells in the cochlea and contribute to K+recycling in the inner ear. Important for the survival of cochlear outer and inner hair cells and the maintenance of the organ of Corti. May be required for basolateral Cl-extrusion in the kidney and contribute

to renal acidification (By similarity).

Synonyms: DKFZP434F076; KCC4




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**




Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA372837 (SLC12A7 Antibody) at dilution 1/35 (Original magnification: ×200)




Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA372837 (SLC12A7 Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification: ×200)





Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA372837 (SLC12A7 Antibody) at dilution 1/35 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA372837 (SLC12A7 Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification: ×200)