

Product datasheet for TA372859

KIR2DL5A Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 1000-5000

WB positive control: Human plasma solution

IHC: 30-150

Positive control: Human liver cancer Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide of human KIR2DL5A Formulation: pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 41 kDa

Gene Name: killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 5A

Database Link: Entrez Gene 57292 Human

Q8N109



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several framework genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response.

Synonyms:

CD158F; KIR2DL5; KIR2DL5.1; KIR2DL5.3; KIR2DL5B

Product images:



Gel: 8%SDS-PAGE Lysate: 40 μg

Lane: Human plasma solution

Primary antibody: TA372859 (KIR2DL5A Antibody)


at dilution 1/800

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 1 minute





Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA372859 (KIR2DL5A Antibody) at dilution 1/35 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TA372859 (KIR2DL5A Antibody) at dilution 1/35, treated with synthetic peptide. (Original magnification: ×200)