

# **Product datasheet for TA372935**

### **HEXB Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB: 500-2000

WB positive control: A172 cell lysate

IHC: 50-100

Positive control: Human tonsil Predicted cell location: Cytoplasm

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide of human HEXB

**Formulation:** pH7.4 PBS, 0.05% NaN3, 40% Glycerol

**Concentration:** lot specific

**Purification:** Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 63 kDa

Gene Name: hexosaminidase subunit beta

Database Link: Entrez Gene 3074 Human

P07686

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






#### Background:

Hexosaminidase B is the beta subunit of the lysosomal enzyme beta-hexosaminidase that, together with the cofactor GM2 activator protein, catalyzes the degradation of the ganglioside GM2, and other molecules containing terminal N-acetyl hexosamines. Beta-hexosaminidase is composed of two subunits, alpha and beta, which are encoded by separate genes. Both beta-hexosaminidase alpha and beta subunits are members of family 20 of glycosyl hydrolases. Mutations in the alpha or beta subunit genes lead to an accumulation of GM2 ganglioside in neurons and neurodegenerative disorders termed the GM2 gangliosidoses. Beta subunit gene mutations lead to Sandhoff disease (GM2-gangliosidosis type II). Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Synonyms:

ENC-1AS; HCC-7; N-acetyl-beta-glucosaminidase

## **Product images:**



Gel: 8%SDS-PAGE Lysate: 40 µg Lane: A172 cell lysate


Primary antibody: TA372935 (HEXB Antibody) at

dilution 1/400

Secondary antibody: Goat anti rabbit IgG at


1/5000 dilution

Exposure time: 3 seconds




Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA372935 (HEXB Antibody) at dilution 1/50 (Original magnification: ×200)






Immunohistochemistry of paraffin-embedded Human tonsil tissue using TA372935 (HEXB Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA372935 (HEXB Antibody) at dilution 1/50 (Original magnification: ×200)



Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using TA372935 (HEXB Antibody) at dilution 1/50, treated with synthetic peptide. (Original magnification: ×200)