

Product datasheet for TA373056S

OriGene Technologies, Inc.9620 Medical Center Drive, Ste 200

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Heparan Sulfate Proteoglycan 2 (HSPG2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 200-1000

WB positive control: LOVO cell lysate

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen:Synthetic peptide of human HSPG2Formulation:pH7.4 PBS, 0.05% NaN3, 40% Glycerol

Concentration: lot specific

Purification: Antigen affinity purification

Conjugation: Unconjugated Storage: Store at -20°C.

Stability: 1 year Predicted Protein Size: 469 kDa

Gene Name: heparan sulfate proteoglycan 2

Database Link: Entrez Gene 3339 Human

P98160




Background:

This gene encodes the perlecan protein, which consists of a core protein to which three long chains of glycosaminoglycans (heparan sulfate or chondroitin sulfate) are attached. The perlecan protein is a large multidomain proteoglycan that binds to and cross-links many extracellular matrix components and cell-surface molecules. It has been shown that this protein interacts with laminin, prolargin, collagen type IV, FGFBP1, FBLN2, FGF7 and transthyretin, etc., and it plays essential roles in multiple biological activities. Perlecan is a key component of the vascular extracellular matrix, where it helps to maintain the endothelial barrier function. It is a potent inhibitor of smooth muscle cell proliferation and is thus thought to help maintain vascular homeostasis. It can also promote growth factor (e.g., FGF2) activity and thus stimulate endothelial growth and re-generation. It is a major component of basement membranes, where it is involved in the stabilization of other molecules as well as being involved with glomerular permeability to macromolecules and cell adhesion. Mutations in this gene cause Schwartz-Jampel syndrome type 1, Silverman-Handmaker type of dyssegmental dysplasia, and tardive dyskinesia. Alternative splicing of this gene results in multiple transcript variants.

Synonyms: HSPG; perlecan; PLC; PRCAN; SJA; SJS; SJS1

Product images:



Gel: 6%SDS-PAGE
Lysate: 40 µg
Lane: LOVO cell lysate
Primary antibody: [TA373056] (HSPG2 Antibody)
at dilution 1/400
Secondary antibody: Goat anti rabbit IgG at

Exposure time: 5 minutes

1/5000 dilution