

Product datasheet for TA373234

ACTR1A Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: WB,1:500 - 1:2000 **Reactivity:** Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 187-376 of

human ACTR1A (NP_005727.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 42kDa

Gene Name: ARP1 actin-related protein 1 homolog A, centractin alpha

Database Link: Entrez Gene 10121 Human

P61163

Background: This gene encodes a 42.6 kD subunit of dynactin, a macromolecular complex consisting of 10-

11 subunits ranging in size from 22 to 150 kD. Dynactin binds to both microtubules and cytoplasmic dynein. It is involved in a diverse array of cellular functions, including ER-to-Golgi

transport, the centripetal movement of lysosomes and endosomes, spindle formation,

chromosome movement, nuclear positioning, and axonogenesis. This subunit is present in 8-

13 copies per dynactin molecule, and is the most abundant molecule in the dynactin

complex. It is an actin-related protein, and is approximately 60% identical at the amino acid

level to conventional actin.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

actin-RPV; ARP1; Centractin; CTRN1; FLJ52695; FLJ52800; FLJ55002

Product images:



Western blot analysis of extracts of various cell lines, using ACTR1A antibody (TA373234) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 5s.