

Product datasheet for TA373253

Primary Antibodies

Human, Mouse, Rat

Recombinant protein of human ADAM15

Store at -20°C. Avoid freeze / thaw cycles.

Shelf life: one year from despatch.

ADAM metallopeptidase domain 15

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

ICC/IF, WB

IF,1:50 - 1:200

Unmodified

Rabbit

Polyclonal

lot specific

Q13444

Affinity purification

55kDa/68kDa/83-92kDa

Entrez Gene 8751 Human

Unconjugated

lgG

ADAM15 Rabbit Polyclonal Antibody

Recommended Dilution: WB.1:500 - 1:2000

Product data:

Product Type:

Applications:

Reactivity:

Host:

Isotype:

Clonality:

Immunogen:

Formulation:

Purification:

Conjugation:

Gene Name: Database Link:

Predicted Protein Size:

Storage:

Stability:

Concentration:

Modifications:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE ADAM15 Rabbit Polyclonal Antibody – TA373253

Background:The protein encoded by this gene is a member of the ADAM (a disintegrin and
metalloproteinase) protein family. ADAM family members are type I transmembrane
glycoproteins known to be involved in cell adhesion and proteolytic ectodomain processing
of cytokines and adhesion molecules. This protein contains multiple functional domains
including a zinc-binding metalloprotease domain, a disintegrin-like domain, as well as a EGF-
like domain. Through its disintegrin-like domain, this protein specifically interacts with the
integrin beta chain, beta 3. It also interacts with Src family protein-tyrosine kinases in a
phosphorylation-dependent manner, suggesting that this protein may function in cell-cell
adhesion as well as in cellular signaling. Multiple alternatively spliced transcript variants
encoding distinct isoforms have been observed.


Synonyms:

MDC-15; MDC15; metargidin

Product images:



Western blot analysis of extracts of various cell lines, using ADAM15 antibody (TA373253) at 1:500 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit . |Exposure time: 90s.



Immunofluorescence analysis of HeLa cells using ADAM15 Polyclonal Antibody (TA373253) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US