

### **Product datasheet for TA373295**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Alcohol Dehydrogenase (ADH1A) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Human, Rat
Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 220-300 of

human Alcohol Dehydrogenase (NP\_000658.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 39kDa

**Gene Name:** alcohol dehydrogenase 1A (class I), alpha polypeptide

Database Link: Entrez Gene 124 Human

P07327

**Background:** This gene encodes a member of the alcohol dehydrogenase family. The encoded protein is

the alpha subunit of class I alcohol dehydrogenase, which consists of several homo- and heterodimers of alpha, beta and gamma subunits. Alcohol dehydrogenases catalyze the oxidation of alcohols to aldehydes. This gene is active in the liver in early fetal life but only

weakly active in adult liver. This gene is found in a cluster with six additional alcohol

dehydrogenase genes, including those encoding the beta and gamma subunits, on the long

arm of chromosome 4. Mutations in this gene may contribute to variation in certain


personality traits and substance dependence.





Synonyms: ADH1

# **Product images:**



Western blot analysis of extracts of various cell lines, using Alcohol Dehydrogenase antibody (TA373295) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.