

# **Product datasheet for TA373413**

## **AKT1 Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

Applications: IP, WB

Recommended Dilution: WB,1:500 - 1:2000

IP,1:20 - 1:50

Reactivity: Human, Mouse, Rat, Zebrafish

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant protein of human AKT1

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 48kDa/55kDa

**Gene Name:** AKT serine/threonine kinase 1

Database Link: Entrez Gene 207 Human

P31749

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn






#### Background:

The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Mutations in this gene have been associated with the Proteus syndrome. Multiple alternatively spliced transcript variants have been found for this gene.

Synonyms:

AKT; C-AKT; MGC99656; PKB; PKB-ALPHA; PRKBA; RAC; RAC-ALPHA; RAC-PK-alpha

### **Product images:**



Western blot analysis of extracts of various cell lines, using AKT1 antibody (TA373413) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.