

## **Product datasheet for TA373433**

## **Albumin (ALB) Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:100 IF,1:50 - 1:100

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant protein of human Albumin

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 45kDa/47kDa/69kDa

Gene Name: albumin

Database Link: Entrez Gene 213 Human

P02768

**Background:** Albumin is a soluble, monomeric protein which comprises about one-half of the blood serum

protein. Albumin functions primarily as a carrier protein for steroids, fatty acids, and thyroid hormones and plays a role in stabilizing extracellular fluid volume. Albumin is a globular unglycosylated serum protein of molecular weight 65,000. Albumin is synthesized in the liver as preproalbumin which has an N-terminal peptide that is removed before the nascent protein is released from the rough endoplasmic reticulum. The product, proalbumin, is in

turn cleaved in the Golgi vesicles to produce the secreted albumin.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

albumin; DKFZp779N1935; PRO0883; PRO0903; PRO1341

## **Product images:**




Western blot analysis of extracts of various cell lines, using Albumin Rabbit pAb (TA373433) at 1:500 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 180s.



Immunofluorescence analysis of HeLa cells using Albumin antibody (TA373433) at dilution of 1:100. Blue: DAPI for nuclear staining.





Immunofluorescence analysis of human liver cancer cells using Albumin antibody (TA373433) at dilution of 1:100. Blue: DAPI for nuclear staining.