

## **Product datasheet for TA373559**

## Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

EU: info-de@origene.com CN: techsupport@origene.cn

## **AP2 alpha (AP2A1) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WE

**Reactivity:** WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of

human AP2A1 (NP\_055018.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 105kDa/107kDa

**Gene Name:** adaptor related protein complex 2 alpha 1 subunit

Database Link: Entrez Gene 160 Human

<u>095782</u>

Background: This gene encodes the alpha 1 adaptin subunit of the adaptor protein 2 (AP-2) complex found

in clathrin coated vesicles. The AP-2 complex is a heterotetramer consisting of two large adaptins (alpha or beta), a medium adaptin (mu), and a small adaptin (sigma). The complex is part of the protein coat on the cytoplasmic face of coated vesicles which links clathrin to receptors in vesicles. Alternative splicing of this gene results in two transcript variants encoding two different isoforms. A third transcript variant has been described, but its full

length nature has not been determined.






Synonyms:

ADTAA; Alpha1-adaptin; AP2-ALPHA; CLAPA1

## **Product images:**



Western blot analysis of extracts of various cell lines, using AP2A1 antibody (TA373559) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.