

## **Product datasheet for TA373630**

## Froduct datasireet for TAS7305

## **ARAP1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

**Reactivity:** Mouse, Rat **Modifications:** Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 834-1133

of human ARAP1 (NP\_001128662.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 77kDa/127kDa/135kDa/136kDa/160kDa/162kDa

**Gene Name:** ArfGAP with RhoGAP domain, ankyrin repeat and PH domain 1

Database Link: Q96P48

**Background:** The protein encoded by this gene contains SAM, ARF-GAP, RHO-GAP, ankyrin repeat, RAS-

associating, and pleckstrin homology (PH) domains. In vitro, this protein displays RHO-GAP and phosphatidylinositol (3,4,5) trisphosphate (PIP3)-dependent ARF-GAP activity. The encoded protein associates with the Golgi, and the ARF-GAP activity mediates changes in the

Golgi and the formation of filopodia. It is thought to regulate the cell-specific trafficking of a receptor protein involved in apoptosis. Multiple transcript variants encoding different

isoforms have been found for this gene.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

Centaurin-delta-2; CENTD2; Cnt-d2; KIAA0782; OTTHUMP00000211462

## **Product images:**



Western blot analysis of extracts of various cell lines, using ARAP1 antibody (TA373630) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.