

## Product datasheet for TA373640

## **ARFGAP1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** 

Recommended Dilution: WB.1:500 - 1:2000 Reactivity: Human, Mouse, Rat

**Modifications:** Unmodified

Host: Rabbit Isotype: **IgG** 

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of

human ARFGAP1 (NP\_001268411.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** 31kDa/39kDa/44kDa/45kDa

Gene Name: ADP ribosylation factor GTPase activating protein 1

Database Link: Entrez Gene 55738 Human

Q8N6T3

Background: The protein encoded by this gene is a GTPase-activating protein, which associates with the

> Golgi apparatus and which interacts with ADP-ribosylation factor 1. The encoded protein promotes hydrolysis of ADP-ribosylation factor 1-bound GTP and is required for the

dissociation of coat proteins from Golgi-derived membranes and vesicles. Dissociation of the

coat proteins is required for the fusion of these vesicles with target compartments. The

activity of this protein is stimulated by phosphoinosides and inhibited by

phosphatidylcholine. Alternative splicing results in multiple transcript variants.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

ARF1GAP; bA261N11.3; FLJ10767; HRIHFB2281; MGC39924

## **Product images:**



Western blot analysis of extracts of various cell lines, using ARFGAP1 antibody (TA373640) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 3s.