

# **Product datasheet for TA373734**

### Asialoglycoprotein Receptor 1 (ASGR1) Rabbit Polyclonal Antibody

### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200 IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

**Host:** Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant protein of human ASGR1

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

**Storage:** Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 29kDa/33kDa

**Gene Name:** asialoglycoprotein receptor 1

Database Link: Entrez Gene 432 Human

P07306



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:


This gene encodes a subunit of the asialoglycoprotein receptor. This receptor is a transmembrane protein that plays a critical role in serum glycoprotein homeostasis by mediating the endocytosis and lysosomal degradation of glycoproteins with exposed terminal galactose or N-acetylgalactosamine residues. The asialoglycoprotein receptor may facilitate hepatic infection by multiple viruses including hepatitis B, and is also a target for liver-specific drug delivery. The asialoglycoprotein receptor is a hetero-oligomeric protein composed of major and minor subunits, which are encoded by different genes. The protein encoded by this gene is the more abundant major subunit. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

Synonyms: ASGPR; CLEC4H1

## **Product images:**




Western blot analysis of extracts of various cell lines, using ASGR1 antibody (TA373734) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.




Immunofluorescence analysis of rat liver using ASGR1 Rabbit pAb (TA373734) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.





Immunofluorescence analysis of human liver cancer using ASGR1 Rabbit pAb (TA373734) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of mouse liver using ASGR1 Rabbit pAb (TA373734) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.