

Product datasheet for TA373737

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Asialoglycoprotein Receptor 2 (ASGR2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Human

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 58-287 of

human ASGR2 (NP_550435.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 32kDa/35kDa

Gene Name: asialoglycoprotein receptor 2

Database Link: Entrez Gene 433 Human

P07307




Background:

This gene encodes a subunit of the asialoglycoprotein receptor. This receptor is a transmembrane protein that plays a critical role in serum glycoprotein homeostasis by mediating the endocytosis and lysosomal degradation of glycoproteins with exposed terminal galactose or N-acetylgalactosamine residues. The asialoglycoprotein receptor may facilitate hepatic infection by multiple viruses including hepatitis B, and is also a target for liver-specific drug delivery. The asialoglycoprotein receptor is a hetero-oligomeric protein composed of major and minor subunits, which are encoded by different genes. The protein encoded by this gene is the less abundant minor subunit. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

Synonyms: ASGP-R; CLEC4H2; L-H2

Product images:



Western blot analysis of extracts of HepG2 cells, using ASGR2 Antibody (TA373737) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.