

Product datasheet for TA373748

ASPA Rabbit Polyclonal Antibody

Product data:

Clonality:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB.1:500 - 1:2000

IF,1:50 - 1:100

Polyclonal

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Rabbit Host: Isotype: lgG

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-313 of

human ASPA (NP_001121557.1).

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. Formulation:

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles. Storage:

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 35kDa

Gene Name: aspartoacylase

Database Link: Entrez Gene 443 Human

P45381

Background: This gene encodes an enzyme that catalyzes the conversion of N-acetyl_L-aspartic acid (NAA)

> to aspartate and acetate. NAA is abundant in the brain where hydrolysis by aspartoacylase is thought to help maintain white matter. This protein is an NAA scavenger in other tissues. Mutations in this gene cause Canavan disease. Alternatively spliced transcript variants have

been found for this gene.

Synonyms: ACY-2; ACY2; Aminoacylase-2; ASP; aspartoacylase




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of extracts of various cell lines, using ASPA Antibody (TA373748) at 1:500 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.



Immunofluorescence analysis of MCF-7 cells using ASPA antibody (TA373748). Blue: DAPI for nuclear staining.