

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA373862

AUH Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 68-339 of human AUH (NP_001689.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	32kDa/35kDa
Gene Name:	AU RNA binding protein/enoyl-CoA hydratase
Database Link:	<u>Entrez Gene 549 Human</u> <u>Q13825</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE AUH Rabbit Polyclonal Antibody – TA373862

AUH

Background:This gene encodes bifunctional mitochondrial protein that has both RNA-binding and
hydratase activities. The encoded protein is a methylglutaconyl-CoA hydratase that catalyzes
the hydration of 3-methylglutaconyl-CoA to 3-hydroxy-3-methyl-glutaryl-CoA, a critical step in
the leucine degradation pathway. This protein also binds AU-rich elements (AREs) found in
the 3' UTRs of rapidly decaying mRNAs including c-fos, c-myc and granulocyte/ macrophage
colony stimulating factor. ARE elements are involved in directing RNA to rapid degradation
and deadenylation. This protein is localizes to the mitochondrial matrix and the inner
mitochondrial membrane and may be involved in mitochondrial protein synthesis. Mutations
in this gene are the cause of 3-methylglutaconic aciduria, type I. Alternative splicing results in
multiple transcript variants.

Synonyms:

Product images:



Western blot analysis of extracts of various cell lines, using AUH antibody (TA373862) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 60s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US