

## **Product datasheet for TA373863**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

## **Aurora A (AURKA) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WE

Recommended Dilution: WB,1:500 - 1:2000

**Reactivity:** Human

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 1-130 of

human AURKA (NP\_940837.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 45kDa

**Gene Name:** aurora kinase A

Database Link: Entrez Gene 6790 Human

<u>014965</u>

**Background:** The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved

in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript variants encoding the

same protein have been found for this gene.






Synonyms:

AIK; ARK1; AURA; Aurora-A; AURORA2; BTAK; hARK1; MGC34538; OTTHUMP00000031344; OTTHUMP00000166071; STK6; STK7; STK15

## **Product images:**



Western blot analysis of extracts of various cell lines, using AURKA antibody (TA373863) at 1:1000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit. |Exposure time: 30s.