

## Product datasheet for **TA373961**

### BCDIN3D Rabbit Polyclonal Antibody


#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000
Reactivity:	Human, Mouse
Modifications:	Unmodified
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-292 of human BCDIN3D (NP_859059.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Gene Name:	BCDIN3 domain containing RNA methyltransferase
Database Link:	<a href="#">Entrez Gene 144233 Human Q7Z5W3</a>
Background:	O-methyltransferase that specifically monomethylates 5'-monophosphate of cytoplasmic histidyl tRNA (tRNA(His, acting as a capping enzyme by protecting tRNA(His from cleavage by DICER1. Also able, with less efficiently, to methylate the 5' monophosphate of a subset of pre-miRNAs, acting as a negative regulator of miRNA processing. The 5' monophosphate of pre-miRNAs is recognized by DICER1 and is required for pre-miRNAs processing: methylation at this position reduces the processing of pre-miRNAs by DICER1. Was also reported to mediate dimethylation of pre-miR-145; however dimethylation cannot be reproduced by another group which observes a monomethylation of pre-miR-145.
Synonyms:	BCDIN3D



[View online »](#)

## Product images:



Western blot analysis of extracts of various cell lines, using BCDIN3D antibody (TA373961) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit. | Exposure time: 180s.