

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA374032

BMP4 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, WB
Recommended Dilution:	WB,1:500 - 1:2000 IF,1:50 - 1:200
Reactivity:	Human, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	lgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-408 of human BMP4 (NP_570912.2).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	46kDa
Gene Name:	bone morphogenetic protein 4
Database Link:	<u>Entrez Gene 652 Human</u> <u>P12644</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE BMP4 Rabbit Polyclonal Antibody – TA374032

Background:This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta)
superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to
recruitment and activation of SMAD family transcription factors that regulate gene
expression. The encoded preproprotein is proteolytically processed to generate each subunit
of the disulfide-linked homodimer. This protein regulates heart development and
adipogenesis. Mutations in this gene are associated with orofacial cleft and microphthalmia
in human patients. The encoded protein may also be involved in the pathology of multiple
cardiovascular diseases and human cancers.

Synonyms: BMP-2B; BMP-4; BMP2B; BMP2B1; DVR4; MCOPS6; OFC11; ZYME

Product images:



Western blot analysis of extracts of A431 cells, using BMP4 antibody (TA374032) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 90s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US