

Product datasheet for TA374191

CACNA2D3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB.1:1000 - 1:2000 Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: **IgG**

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 900-1040

of human CACNA2D3 (NP_060868.2).

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. Formulation:

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 59kDa/112kDa/123kDa

Gene Name: calcium voltage-gated channel auxiliary subunit alpha2delta 3

Database Link: Entrez Gene 55799 Human

Q8IZS8

Background: This gene encodes a member of the alpha-2/delta subunit family, a protein in the voltage-

> dependent calcium channel complex. Calcium channels mediate the influx of calcium ions into the cell upon membrane polarization and consist of a complex of alpha-1, alpha-2/delta, beta, and gamma subunits in a 1:1:1:1 ratio. Various versions of each of these subunits exist, either expressed from similar genes or the result of alternative splicing. Research on a highly similar protein in rabbit suggests the protein described in this record is cleaved into alpha-2 and delta subunits. Alternate transcriptional splice variants of this gene have been observed

but have not been thoroughly characterized.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: HSA272268

Product images:



Western blot analysis of extracts of various cell lines, using CACNA2D3 antibody (TA374191) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.