

Product datasheet for TA374263

i i oddet datasiieet ioi iA5/420.

CARD10 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200 IF,1:50 - 1:200

Polyclonal

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit

Isotype: IgG

Clonality:

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 863-1032

of human CARD10 (NP 055365.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 82kDa/115kDa

Gene Name: caspase recruitment domain family member 10

Database Link: Entrez Gene 29775 Human

Q9BWT7

Background: The caspase recruitment domain (CARD) is a protein module that consists of 6 or 7

antiparallel alpha helices. It participates in apoptosis signaling through highly specific protein-protein homophilic interactions. Like several other CARD proteins, CARD10 belongs to the membrane-associated guanylate kinase (MAGUK) family and activates NF-kappa-B (NFKB; see MIM 164011) through BCL10 (MIM 603517) (Wang et al., 2001 [PubMed 11259443]).[supplied

by OMIM, Mar 2008]



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

BIMP1; CARMA3; MGC142219; OTTHUMP00000197888

Product images:



Western blot analysis of extracts of Mouse kidney, using CARD10 antibody (TA374263) at 1:500 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit. |Exposure time: 10s.



Western blot analysis of extracts of Rat kidney, using CARD10 antibody (TA374263) at 1:500 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: