

# Product datasheet for TA374291S

## FLASH (CASP8AP2) Rabbit Polyclonal Antibody

## **Product data:**

#### **Product Type: Primary Antibodies Applications:** ELISA, WB Recommended Dilution: WB.1:500 - 1:2000 ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. **Reactivity:** Human Modifications: Unmodified Host: Rabbit lgG Isotype: **Clonality:** Polyclonal Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. **Concentration:** lot specific Purification: Affinity purification **Conjugation:** Unconjugated Store at -20°C. Avoid freeze / thaw cycles. Storage: Shelf life: one year from despatch. Stability: **Predicted Protein Size:** 223kDa Gene Name: caspase 8 associated protein 2 Database Link: Entrez Gene 9994 Human Q9UKL3 Background: This protein is highly similar to FLASH, a mouse apoptotic protein identified by its interaction with the death-effector domain (DED) of caspase 8. Studies of FLASH protein suggested that this protein may be a component of the death-inducing signaling complex that includes Fas receptor, Fas-binding adapter FADD, and caspase 8, and plays a regulatory role in Fasmediated apoptosis. Alternative splicing results in multiple transcript variants encoding the same protein. Synonyms: CED-4; FLASH; FLJ11208; KIAA1315; RIP25




View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **Product images:**



Western blot analysis of various lysates using CASP8AP2 Rabbit pAb ([TA374291]) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US