

Product datasheet for TA374314

CBL Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	ICC/IF, IHC, IP, WB
Recommended Dilution:	WB,1:500 - 1:2000 IHC,1:50 - 1:100
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 637-906 of human CBL (NP_005179.2).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	99kDa
Gene Name:	Cbl proto-oncogene
Database Link:	<u>Entrez Gene 867 Human</u> <u>P22681</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE CBL Rabbit Polyclonal Antibody – TA374314

Background:This gene is a proto-oncogene that encodes a RING finger E3 ubiquitin ligase. The encoded
protein is one of the enzymes required for targeting substrates for degradation by the
proteasome. This protein mediates the transfer of ubiquitin from ubiquitin conjugating
enzymes (E2) to specific substrates. This protein also contains an N-terminal phosphotyrosine
binding domain that allows it to interact with numerous tyrosine-phosphorylated substrates
and target them for proteasome degradation. As such it functions as a negative regulator of
many signal transduction pathways. This gene has been found to be mutated or translocated
in many cancers including acute myeloid leukaemia, and expansion of CGG repeats in the 5'
UTR has been associated with Jacobsen syndrome. Mutations in this gene are also the cause
of Noonan syndrome-like disorder.

Synonyms:

C-CBL; CBL2; RNF55

Product images:



Western blot analysis of extracts of various cell lines, using CBL antibody (TA374314) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 90s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US