

Product datasheet for TA374386

Cyclin G (CCNG1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse

Modifications: Unmodified

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-295 of

human Cyclin G1 (NP_004051.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 18kDa/34kDa

Gene Name: cyclin G1

Database Link: Entrez Gene 900 Human

P51959

Background: The eukaryotic cell cycle is governed by cyclin-dependent protein kinases (CDKs) whose

activities are regulated by cyclins and CDK inhibitors. The protein encoded by this gene is a member of the cyclin family and contains the cyclin box. The encoded protein lacks the

protein destabilizing (PEST) sequence that is present in other family members. Transcriptional

activation of this gene can be induced by tumor protein p53. Two transcript variants

encoding the same protein have been identified for this gene.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: CCNG; CYCG1; Cyclin-G

Product images:



Western blot analysis of extracts of mouse skeletal muscle, using Cyclin G1 antibody (TA374386) at 1:1000 dilution.|Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Enhanced Kit.|Exposure time: 90s.