

Product datasheet for TA374623

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

EU: info-de@origene.com
CN: techsupport@origene.cn

Bile salt activated lipase (CEL) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Reactivity: WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic peptide of human CEL

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 71kDa/79kDa

Gene Name: carboxyl ester lipase

Database Link: Entrez Gene 1056 Human

P19835

Background: The protein encoded by this gene is a glycoprotein secreted from the pancreas into the

digestive tract and from the lactating mammary gland into human milk. The physiological role of this protein is in cholesterol and lipid-soluble vitamin ester hydrolysis and absorption. This encoded protein promotes large chylomicron production in the intestine. Also its presence in plasma suggests its interactions with cholesterol and oxidized lipoproteins to modulate the progression of atherosclerosis. In pancreatic tumoral cells, this encoded protein is thought to be sequestrated within the Golgi compartment and is probably not secreted. This gene

contains a variable number of tandem repeat (VNTR) polymorphism in the coding region that

may influence the function of the encoded protein.






Synonyms:

BAL; BSDL; BSSL; Bucelipase; CEase; CELL; FAP; FAPP; LIPA; MODY8

Product images:



Western blot analysis of extracts of various cell lines, using CEL antibody (TA374623) at 1:500 dilution._Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution._Lysates/proteins: 25ug per lane._Blocking buffer: 3% nonfat dry milk in TBST._Detection: ECL Enhanced Kit ._Exposure time: 30s.