

## **Product datasheet for TA374788**

## **CHST15 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** 

Recommended Dilution: WB.1:500 - 1:2000

Reactivity: Human, Rat **Modifications:** Unmodified

Host: Rabbit Isotype: **IgG** 

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 452-561 of

human CHST15 (NP\_056976.2).

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. Formulation:

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** 58kDa/64kDa

Gene Name: carbohydrate (N-acetylgalactosamine 4-sulfate 6-O) sulfotransferase 15

**Database Link:** Entrez Gene 51363 Human

Q7LFX5

Background: Chondroitin sulfate (CS) is a glycosaminoglycan which is an important structural component

> of the extracellular matrix and which links to proteins to form proteoglycans. Chondroitin sulfate E (CS-E) is an isomer of chondroitin sulfate in which the C-4 and C-6 hydroxyl groups are sulfated. This gene encodes a type II transmembrane glycoprotein that acts as a

> sulfotransferase to transfer sulfate to the C-6 hydroxal group of chondroitin sulfate. This gene has also been identified as being co-expressed with RAG1 in B-cells and as potentially acting as a B-cell surface signaling receptor. Alternative splicing results in multiple transcript

variants encoding distinct isoforms.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

BRAG; DKFZp781H1369; GALNAC4S-6ST; GALNAC4S6ST; hBRAG; KIAA0598; MGC34346; OTTHUMP00000020689; RP11-47G11.1

## **Product images:**



Western blot analysis of extracts of various cell lines, using CHST15 antibody (TA374788) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.