

## Product datasheet for **TA374800**

### CIAO1 Rabbit Polyclonal Antibody

#### Product data:


|                         |   |
|-------------------------|---|
| Product Type:           | Primary Antibodies  |
| Applications:           | WB  |
| Recommended Dilution:   | WB,1:500 - 1:2000   |
| Reactivity:             | Human, Mouse, Rat   |
| Modifications:          | Unmodified  |
| Host:                   | Rabbit  |
| Isotype:                | IgG   |
| Clonality:              | Polyclonal  |
| Immunogen:              | Recombinant fusion protein containing a sequence corresponding to amino acids 1-339 of human CIAO1 (NP_004795.1).   |
| Formulation:            | Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.   |
| Concentration:          | lot specific  |
| Purification:           | Affinity purification   |
| Conjugation:            | Unconjugated  |
| Storage:                | Store at -20°C. Avoid freeze / thaw cycles.   |
| Stability:              | Shelf life: one year from despatch.   |
| Predicted Protein Size: | 37kDa   |
| Gene Name:              | cytosolic iron-sulfur assembly component 1  |
| Database Link:          | <a href="#">Entrez Gene 9391 Human O76071</a>   |
| Background:             | Key component of the cytosolic iron-sulfur protein assembly (CIA complex, a multiprotein complex that mediates the incorporation of iron-sulfur cluster into extramitochondrial Fe/S proteins. As a CIA complex component, interacts specifically with CIAO2A or CIAO2B and MMS19 to assist different branches of iron-sulfur protein assembly, depending of its interactors. The complex CIAO1:CIAO2B:MMS19 binds to and facilitates the assembly of most cytosolic-nuclear Fe/S proteins. CIAO1:CIAO2A specifically matures ACO1 and stabilizes IREB2. Seems to specifically modulate the transactivation activity of WT1. As part of the mitotic spindle-associated MMXD complex it may play a role in chromosome segregation. |



[View online »](#)

Synonyms: CIA1; WDR39

### Product images:



Western blot analysis of extracts of various cell lines, using CIAO1 antibody (TA374800) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit . | Exposure time: 90s.