

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA374881

CLK2 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 239-498 of human CLK2 (NP_003984.2).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	17kDa/59kDa/60kDa
Gene Name:	CDC like kinase 2
Database Link:	<u>Entrez Gene 1196 Human</u> <u>P49760</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE CLK2 Rabbit Polyclonal Antibody – TA374881

Background:This gene encodes a dual specificity protein kinase that phosphorylates serine/threonine and
tyrosine-containing substrates. Activity of this protein regulates serine- and arginine-rich (SR)
proteins of the spliceosomal complex, thereby influencing alternative transcript splicing.
Chromosomal translocations have been characterized between this locus and the PAFAH1B3
(platelet-activating factor acetylhydrolase 1b, catalytic subunit 3 (29kDa)) gene on
chromosome 19, resulting in the production of a fusion protein. Note that this gene is distinct
from the TELO2 gene (GeneID:9894), which shares the CLK2 alias, but encodes a protein that
is involved in telomere length regulation. There is a pseudogene for this gene on
chromosome 7. Alternative splicing results in multiple transcript variants.

Synonyms:

hCLK2; MGC61500

Product images:



Western blot analysis of extracts of various cell lines, using CLK2 antibody (TA374881) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 30s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US