

## **Product datasheet for TA374913**

## Product datasireet for TAS/4913

## Product data:

**Product Type:** Primary Antibodies

**CNBP Rabbit Polyclonal Antibody** 

**Applications:** ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse

Modifications: Unmodified

**Host:** Rabbit

**Isotype:** IgG

Clonality: Polyclonal

**Immunogen:** A synthetic peptide corresponding to a sequence within amino acids 100 to the C-terminus of

human CNBP (NP\_001120668.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 17kDa/18kDa/19kDa

**Gene Name:** CCHC-type zinc finger nucleic acid binding protein

Database Link: Entrez Gene 7555 Human

P62633

**Background:** This gene encodes a nucleic-acid binding protein with seven zinc-finger domains. The protein

has a preference for binding single stranded DNA and RNA. The protein functions in capindependent translation of ornithine decarboxylase mRNA, and may also function in sterolmediated transcriptional regulation. A CCTG expansion from <30 repeats to 75-11000 repeats

in the first intron of this gene results in myotonic dystrophy type 2. Multiple transcript

variants encoding different isoforms have been found for this gene.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

CNBP1; DM2; FLJ11631; PROMM; RNF163; ZCCHC22; ZNF9

## **Product images:**



Western blot analysis of extracts of various cell lines, using CNBP antibody (TA374913) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 5s.



Immunofluorescence analysis of U-2 OS cells using CNBP antibody (TA374913) at dilution of 1:100. Blue: DAPI for nuclear staining.