

## **Product datasheet for TA374973**

## Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

EU: info-de@origene.com CN: techsupport@origene.cn

## Collagen IX (COL9A1) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 20-270 of

human COL9A1 (NP\_001842.3).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 35kDa/64kDa/91kDa

**Gene Name:** collagen type IX alpha 1

Database Link: P20849

**Background:** This gene encodes one of the three alpha chains of type IX collagen, which is a minor (5-20%)

collagen component of hyaline cartilage. Type IX collagen is usually found in tissues containing type II collagen, a fibrillar collagen. Studies in knockout mice have shown that synthesis of the alpha 1 chain is essential for assembly of type IX collagen molecules, a heterotrimeric molecule, and that lack of type IX collagen is associated with early onset osteoarthritis. Mutations in this gene are associated with osteoarthritis in humans, with multiple epiphyseal dysplasia, 6, a form of chondrodysplasia, and with Stickler syndrome, a disease characterized by ophthalmic, orofacial, articular, and auditory defects. Two transcript

variants that encode different isoforms have been identified for this gene.






Synonyms:

DJ149L1.1.2; EDM6; FLJ40263; MED

## **Product images:**



Western blot analysis of extracts of mouse skeletal muscle, using COL9A1 antibody (TA374973) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.