

Product datasheet for TA375015

COX5A Rabbit Polyclonal Antibody

Product data:

Isotype:

Product Type: Primary Antibodies

Applications: ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:100 IF,1:10 - 1:100

Reactivity: Human, Mouse, Rat

IgG

Modifications: Unmodified

Host: Rabbit

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-150 of

human COX5A (NP_004246.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 16kDa

Gene Name: cytochrome c oxidase subunit 5A

Database Link: Entrez Gene 9377 Human

P20674



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:


Cytochrome c oxidase (COX) is the terminal enzyme of the mitochondrial respiratory chain. It is a multi-subunit enzyme complex that couples the transfer of electrons from cytochrome c to molecular oxygen and contributes to a proton electrochemical gradient across the inner mitochondrial membrane. The complex consists of 13 mitochondrial- and nuclear-encoded subunits. The mitochondrially-encoded subunits perform the electron transfer of proton pumping activities. The functions of the nuclear-encoded subunits are unknown but they may play a role in the regulation and assembly of the complex. This gene encodes the nuclear-encoded subunit Va of the human mitochondrial respiratory chain enzyme. A pseudogene COX5AP1 has been found in chromosome 14q22.

Synonyms: COX; COX-VA; VA

Product images:



Western blot analysis of extracts of various cell lines, using COX5A Antibody (TA375015) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.



Immunofluorescence analysis of U2OS cells using COX5A antibody (TA375015). Blue: DAPI for nuclear staining.