

Product datasheet for TA375020

COX7A1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200

Reactivity: Mouse, Rat, Human

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-79 of

human COX7A1 (NP_001855.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 9kDa

Gene Name: cytochrome c oxidase subunit 7A1

Database Link: Entrez Gene 1346 Human

P24310



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes polypeptide 1 (muscle isoform) of subunit VIIa and the polypeptide 1 is present only in muscle tissues. Other polypeptides of subunit VIIa are present in both muscle and nonmuscle tissues, and are encoded by different genes.

Synonyms: COX7A; COX7AH; COX7AM

Product images:



Western blot analysis of extracts of Mouse heart, using COX7A1 antibody (TA375020) at 1:500 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.