

Product datasheet for TA375064S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CRABP2 Rabbit Polyclonal Antibody

Product data:

Applications:

Product Type: Primary Antibodies

Recommended Dilution: WB.1:500 - 1:2000

IF/ICC,1:20 - 1:100

ELISA, ICC/IF, WB

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 16kDa

Gene Name: cellular retinoic acid binding protein 2

Database Link: Entrez Gene 1382 Human

P29373

Background: This gene encodes a member of the retinoic acid (RA, a form of vitamin A) binding protein

family and lipocalin/cytosolic fatty-acid binding protein family. The protein is a cytosol-to-nuclear shuttling protein, which facilitates RA binding to its cognate receptor complex and transfer to the nucleus. It is involved in the retinoid signaling pathway, and is associated with increased circulating low-density lipoprotein cholesterol. Alternatively spliced transcript

variants encoding the same protein have been found for this gene.


variants encouning the same protein have been round for this g

Synonyms: CRABP-II; OTTHUMP00000038732; RBP6





Product images:



Western blot analysis of various lysates using CRABP2 Rabbit pAb ([TA375064]) at 1:1000 dilution._Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1:10000 dilution._Lysates/proteins: 25µg per lane._Blocking buffer: 3% nonfat dry milk in TBST._Detection: ECL Enhanced Kit ._Exposure time: 30s.

Immunofluorescence analysis of U2OS cells using CRABP2 Rabbit pAb ([TA375064]). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.