

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA375111

TORC2 (CRTC2) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000
Reactivity:	Human
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 434-693 of human CRTC2 (NP_859066.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	73kDa
Gene Name:	CREB regulated transcription coactivator 2
Database Link:	<u>Entrez Gene 200186 Human</u> <u>Q53ET0</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE TORC2 (CRTC2) Rabbit Polyclonal Antibody – TA375111

Background:This gene encodes a member of the transducers of regulated cAMP response element-
binding protein activity family of transcription coactivators. These proteins promote the
transcription of genes targeted by the cAMP response element-binding protein, and
therefore play an important role in many cellular processes. Under basal conditions the
encoded protein is phosphorylated by AMP-activated protein kinase or the salt-inducible
kinases and is sequestered in the cytoplasm. Upon activation by elevated cAMP or calcium,
the encoded protein translocates to the nucleus and increases target gene expression. Single
nucleotide polymorphisms in this gene may increase the risk of type 2 diabetes. A
pseudogene of this gene is located on the long arm of chromosome 5.

Synonyms:

RP11-422P24.6; TORC-2; TORC2

Product images:



Western blot analysis of extracts of various cell lines, using CRTC2 antibody (TA375111) at 1:500 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit . | Exposure time: 180s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US