

# Product datasheet for TA375165S

## Cystatin SN (CST1) Rabbit Polyclonal Antibody

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	WB,1:500 - 1:2000 ELISA,Recommended starting concentration is 1 $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements.
Reactivity:	Human
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	14kDa
Gene Name:	cystatin SN
Database Link:	<u>Entrez Gene 1469 Human</u> <u>P01037</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **GRIGENE** Cystatin SN (CST1) Rabbit Polyclonal Antibody – TA375165S

Background:The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences.<br/>Some of the members are active cysteine protease inhibitors, while others have lost or<br/>perhaps never acquired this inhibitory activity. There are three inhibitory families in the<br/>superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The<br/>type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of<br/>human fluids and secretions, where they appear to provide protective functions. The cystatin<br/>locus on chromosome 20 contains the majority of the type 2 cystatin genes and<br/>pseudogenes. This gene is located in the cystatin locus and encodes a cysteine proteinase<br/>inhibitor found in saliva, tears, urine, and seminal fluid.

Synonyms: Cystain-SA-I; Cystatin-1; OTTHUMP00000164184

#### **Product images:**



Western blot analysis of various lysates using CST1 Rabbit pAb ([TA375165]) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US