

## **Product datasheet for TA375182S**

## **CTBP2 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A synthetic peptide corresponding to a sequence within amino acids 300 to the C-terminus of

human CTBP2 (NP\_001320.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 48kDa/56kDa/106kDa

Gene Name: C-terminal binding protein 2

Database Link: Entrez Gene 1488 Human

P56545

**Background:** This gene produces alternative transcripts encoding two distinct proteins. One protein is a

transcriptional repressor, while the other isoform is a major component of specialized

synapses known as synaptic ribbons. Both proteins contain a NAD+ binding domain similar to NAD+-dependent 2-hydroxyacid dehydrogenases. A portion of the 3' untranslated region was

used to map this gene to chromosome 21q21.3; however, it was noted that similar loci elsewhere in the genome are likely. Blast analysis shows that this gene is present on chromosome 10. Several transcript variants encoding two different isoforms have been

found for this gene.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: CTBP2


## **Product images:**



Western blot analysis of extracts of various cell lines, using CTBP2 antibody ([TA375182]) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s

Immunofluorescence analysis of C6 cells using CTBP2 antibody ([TA375182]) at dilution of 1:100. Blue: DAPI for nuclear staining.





Immunofluorescence analysis of U2OS cells using CTBP2 antibody ([TA375182]) at dilution of 1:100. Blue: DAPI for nuclear staining.