

## **Product datasheet for TA375203**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### beta Catenin (CTNNB1) Rabbit Polyclonal Antibody

### **Product data:**

**Product Type:** Primary Antibodies

Applications: WE

**Reactivity:** WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Phospho S45

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic phosphorylated peptide around S45 of human CTNNB1 (NP\_001091679).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 9kDa/85kDa **Gene Name:** catenin beta 1

Database Link: Entrez Gene 1499 Human

P35222

**Background:** The protein encoded by this gene is part of a complex of proteins that constitute adherens

junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon.

Mutations in this gene are a cause of colorectal cancer (CRC), pilomatrixoma (PTR),

medulloblastoma (MDB), and ovarian cancer. Alternative splicing results in multiple transcript

variants.






Synonyms:

beta-catenin; CTNNB; DKFZp686D02253; FLJ25606; FLJ37923; OTTHUMP00000209288

# **Product images:**



Western blot analysis of extracts of HeLa cells, using Phospho-β-Catenin-S45 antibody (TA375203) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% BSA.