

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA375282

CYP27B1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000
Reactivity:	Human, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 25-195 of human CYP27B1 (NP_000776.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	56kDa
Gene Name:	cytochrome P450 family 27 subfamily B member 1
Database Link:	<u>Entrez Gene 1594 Human</u> <u>O15528</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE CYP27B1 Rabbit Polyclonal Antibody – TA375282

Background:This gene encodes a member of the cytochrome P450 superfamily of enzymes. The
cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in
drug metabolism and synthesis of cholesterol, steroids and other lipids. The protein encoded
by this gene localizes to the inner mitochondrial membrane where it hydroxylates 25-
hydroxyvitamin D3 at the 1alpha position. This reaction synthesizes 1alpha,25-
dihydroxyvitamin D3, the active form of vitamin D3, which binds to the vitamin D receptor
and regulates calcium metabolism. Thus this enzyme regulates the level of biologically active
vitamin D and plays an important role in calcium homeostasis. Mutations in this gene can
result in vitamin D-dependent rickets type I.

Synonyms: 1alpha(OH)ase; CP2B; CYP1; CYP1alpha; CYP27B; P450c1; P450C1-alpha; P450VD1-alpha; PDDR; VDD1; VDDR; VDDR; VDDR]

Product images:



Western blot analysis of extracts of various cell lines, using CYP27B1 antibody (TA375282) at 1:3000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit . |Exposure time: 90s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US