

## **Product datasheet for TA375329S**

## DAZAP2 Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

**Reactivity:** Mouse

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 1-168 of

human DAZAP2 (NP\_055579.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 9kDa/13kDa/15kDa/16kDa/17kDa/22kDa

**Gene Name:** DAZ associated protein 2

Database Link: Q15038

**Background:** This gene encodes a proline-rich protein which interacts with the deleted in azoospermia

(DAZ) and the deleted in azoospermia-like gene through the DAZ-like repeats. This protein also interacts with the transforming growth factor-beta signaling molecule SARA (Smad anchor for receptor activation), eukaryotic initiation factor 4G, and an E3 ubiquitinase that regulates its stability in splicing factor containing nuclear speckles. The encoded protein may function in various biological and pathological processes including spermatogenesis, cell signaling and transcription regulation, formation of stress granules during translation arrest, RNA splicing, and pathogenesis of multiple myeloma. Multiple transcript variants encoding

different isoforms have been found for this gene.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

KIAA0058; MGC766; MGC14319; PRTB

## **Product images:**



Western blot analysis of extracts of various cell lines, using DAZAP2 Rabbit pAb ([TA375329]) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.