

Product datasheet for TA375331S

DAZL Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200

IF,1:50 - 1:200

Reactivity: Mouse, Rat **Modifications:** Unmodified

Host: Rabbit

Isotype: lgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 186-295 of

human DAZL (NP_001342.2).

Formulation: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 33kDa/35kDa

Gene Name: deleted in azoospermia like

Database Link: Q92904



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The DAZ (Deleted in AZoospermia) gene family encodes potential RNA binding proteins that are expressed in prenatal and postnatal germ cells of males and females. The protein encoded by this gene is localized to the nucleus and cytoplasm of fetal germ cells and to the cytoplasm of developing oocytes. In the testis, this protein is localized to the nucleus of spermatogonia but relocates to the cytoplasm during meiosis where it persists in spermatids and spermatozoa. Transposition and amplification of this autosomal gene during primate evolution gave rise to the DAZ gene cluster on the Y chromosome. Mutations in this gene have been linked to severe spermatogenic failure and infertility in males. Two transcript variants encoding different isoforms have been found for this gene.


Synonyms:

DAZH; DAZL1; DAZLA; MGC26406; SPGY-like-autosomal; SPGYLA

Product images:




Western blot analysis of extracts of various cell lines, using DAZL antibody ([TA375331]) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.



Immunofluorescence analysis of rat testis using DAZL Rabbit pAb ([TA375331]) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.





Immunofluorescence analysis of mouse testis using DAZL Rabbit pAb ([TA375331]) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.