

Product datasheet for **TA375355**

DCPS Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:1000 - 1:2000
Reactivity:	Human
Modifications:	Unmodified
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-270 of human DCPS (NP_054745.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	38kDa
Gene Name:	decapping enzyme, scavenger
Database Link:	Entrez Gene 28960 Human Q96C86




[View online »](#)

Background:

Decapping scavenger enzyme that catalyzes the cleavage of a residual cap structure following the degradation of mRNAs by the 3'→5' exosome-mediated mRNA decay pathway. Hydrolyzes cap analog structures like 7-methylguanosine nucleoside triphosphate (m7GpppG) with up to 10 nucleotide substrates (small capped oligoribonucleotides and specifically releases 5'-phosphorylated RNA fragments and 7-methylguanosine monophosphate (m7GMP. Cleaves cap analog structures like tri-methyl guanosine nucleoside triphosphate (m3(2,2,7GpppG with very poor efficiency. Does not hydrolyze unmethylated cap analog (GpppG and shows no decapping activity on intact m7GpppG-capped mRNA molecules longer than 25 nucleotides. Does not hydrolyze 7-methylguanosine diphosphate (m7GDP to m7GMP. May also play a role in the 5'→3' mRNA decay pathway; m7GDP, the downstream product released by the 5'→3' mRNA mediated decapping activity, may be also converted by DCPS to m7GMP. Binds to m7GpppG and strongly to m7GDP. Plays a role in first intron splicing of pre-mRNAs. Inhibits activation-induced cell death.

Synonyms:

DCS-1; DCS1; HINT-5; HINT5; HSL1; HSPC015

Product images:

Western blot analysis of extracts of HT-29 cells, using DCPS antibody (TA375355) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 15s.