

Product datasheet for TA375363

DCUN1D1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB.1:500 - 1:2000

Reactivity: Human, Mouse

Modifications: Unmodified

Host: Rabbit

Isotype: **IgG**

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 15-70 of

human DCUN1D1 (NP_065691.2).

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. Formulation:

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 30kDa

Gene Name: defective in cullin neddylation 1 domain containing 1

Database Link: Entrez Gene 54165 Human

Q96GG9

Background: Part of an E3 ubiquitin ligase complex for neddylation. Promotes neddylation of cullin

> components of E3 cullin-RING ubiquitin ligase complexes. Acts by binding to cullin-RBX1 complexes in the cytoplasm and promoting their nuclear translocation, enhancing recruitment of E2-NEDD8 (UBE2M-NEDD8 thioester to the complex, and optimizing the orientation of proteins in the complex to allow efficient transfer of NEDD8 from the E2 to the

> cullin substrates. Involved in the release of inhibitory effets of CAND1 on cullin-RING ligase E3

complex assembly and activity. Acts also as an oncogene facilitating malignant

transformation and carcinogenic progression (By similarity.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

DCUN1L1; RP42; SCCRO; SCRO; Tes3

Product images:



Western blot analysis of extracts of various cell lines, using DCUN1D1 antibody (TA375363) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.