

Product datasheet for **TA375378**

MCK10 (DDR1) Rabbit Polyclonal Antibody

Product data:


Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000
Reactivity:	Human, Mouse, Rat
Modifications:	Phospho Y792
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	A phospho synthetic peptide corresponding to residues surrounding Y792 of human DDR1.
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	27kDa/97kDa/99kDa/101kDa
Gene Name:	discoidin domain receptor tyrosine kinase 1
Database Link:	Entrez Gene 780 Human Q08345
Background:	Receptor tyrosine kinases play a key role in the communication of cells with their microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. The protein encoded by this gene belongs to a subfamily of tyrosine kinase receptors with homology to Dictyostelium discoideum protein discoidin I in their extracellular domain, and that are activated by various types of collagen. Expression of this protein is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Feb 2011]



[View online »](#)

Synonyms: CAK; CD167; DDR; EDDR1; HGK2; MCK-10; MCK10; NEP; NTRK4; OTTHUMP00000029344; OTTHUMP00000029345; OTTHUMP00000029346; OTTHUMP00000029347; PTK3; PTK3A; RTK6; TRKE

Product images:



Western blot analysis of extracts of various cell lines, using Phospho-DDR1-Y792 antibody (TA375378) at 1:1000 dilution. 293T cells were treated by ATP(5 mM) at 30°C for 1 hour. NIH/3T3 cells were treated by ATP(5 mM) at 30°C for 1 hour. C6 cells were treated by ATP(5 mM) at 30°C for 1 hour. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% BSA. |Detection: ECL Basic Kit . |Exposure time: 1s.